

AS1484 (Revised October 2024)

# Harvesting, Storing and Feeding High-moisture Corn

Reviewed by

**Karl Hoppe**, Extension Livestock Systems Specialist, Carrington Research Extension Center  
**Colin Tobin**, Animal Scientist, Carrington Research Extension Center

In years with cool summer temperatures and above-average moisture many producers exploring alternative harvesting options for high-moisture grain corn. For livestock and nearby grain producers, harvesting corn as high-moisture grain is one option that will eliminate grain-drying costs and produce a product that makes excellent feed for ruminant animals.

High-moisture corn (HMC) offers many advantages for producers who feed beef or dairy cattle. However, successfully using high-moisture corn requires attention to harvest timing, processing, storage conditions and feeding management.

## What is High-moisture Corn?

High-moisture corn refers to corn harvested at 24% or greater moisture, stored and allowed to ferment in a silo or other storage structure, and used as feed for livestock.

## Advantages of High-moisture Corn

- High-moisture corn is similar in energy and protein content to dry corn.
- No grain drying costs are encountered.
- Yields typically are increased due to less ear drop in the field.
- Longer-season (greater relative maturity) corn can be grown with greater yield potential.
- High-moisture corn allows for an earlier corn harvest, avoiding difficulties associated with adverse weather and potentially spreading the workload during harvest.
- Cattle have a longer window for grazing corn residue.

## Disadvantages of High-moisture Corn

- Marketing alternatives are limited. High-moisture corn is marketable only through livestock. Since it has fermented, high-moisture corn cannot be marketed for ethanol production or other uses typical for dry corn.
- High-moisture corn may require additional storage and processing equipment.
- Improperly ensiling high-moisture corn will result in excessive spoilage and storage losses.
- High-moisture corn tends to ferment faster in the rumen and requires better bunk management, compared with dry corn.

## Harvesting

High-moisture corn should be harvested at 24% to 33% moisture for optimum storage (maximum moisture of 40 percent). Harvesting corn at these moisture levels may require adjustments to harvest equipment to ensure grain is removed from the cob uniformly. Some damage to the grain is acceptable because it still will need to be processed prior to ensiling.

## Processing

High-moisture corn should be processed (ground or rolled) prior to storage. Grinding or rolling and subsequent packing of the corn facilitates oxygen exclusion in the silo. The goal with processing should be to have no more than 5% fines and no more than 5% whole kernels.

For feeding purposes, rolling is the preferred method of processing (compared with grinding) because rolling will result in less fines and lower probability of acidosis due to rapid fermentation when feeding.

**Photo 1** shows a high-capacity hammermill specially designed for high-volume processing of high-moisture corn.

**Photo 2** shows a tub grinder adapted to grind corn into a bunker silo.

While grinding may lead to faster ruminal fermentation and greater risk of acidosis, this problem is an issue only when feeding rations that contain greater than 50% HMC. With the increased feeding of wet distillers grains, the risk of acidosis from feeding ground high-moisture corn is greatly reduced due to the reduction in total ration starch content. Likewise, most dairies finely grind the high-moisture corn to facilitate greater ruminal digestion, and since grain is generally less than 50% of the diet, acidosis is generally not a problem.


## Storage

At moisture levels below 27%, water may need to be added to facilitate packing and fermentation. A good rule of thumb is to add 3.5 gallons of water per ton of corn for each percentage point the corn is below 27% moisture.



**Photo 1. High-moisture grain being processed with high-volume hammermill for storage in a bunker silo.**

*(photo courtesy of DuraTech Industries International, Inc.)*



**Photo 2. High-moisture grain being processed with a tub grinder.**

*(photo courtesy of DuraTech Industries International, Inc.)*

## Bunker or Trench Silo

A bunker or trench silo is the best option for large volumes of corn harvested in a short period of time but will require a large-volume roller or hammermill to process grain rapidly. Packing can be done with a tractor equipped with a dozer blade or a front-end loader. During feedout, the face of the bunker must be kept fresh to avoid

heating, so the width of the high-moisture corn bunker should match the rate of use. Multiple narrower bunkers may be more useful than one wide bunker silo. Bunkers constructed of a cement base and sides will reduce spoilage better than earthen structures and provide a firm surface for equipment throughout the year.



Proper packing of the high-moisture corn in bunker silos also is critical to oxygen exclusion. **Photo 3** shows a properly packed, covered and faced bunker silo filled with high-moisture corn. Proper packing, covering and facing will reduce spoilage losses and improve acceptability of the high-moisture corn in the feed bunk.

A minimum density of 45 pounds of dry matter per cubic foot is desirable for good-quality high-moisture corn. Packing grain in bunker silos can be dangerous, so be sure an experienced and safe tractor operator is operating the packing equipment.

Also, cover all bunker, trench and pile structures with plastic to create an anaerobic environment and minimize spoilage. Inspect plastic covers periodically, and repair any holes or tears.

## Plastic Silage Bags

**Photo 4** shows a large-capacity bagging machine filling a plastic silage bag with high-moisture grain. Bags are filled with a specialized bagger that also will process the corn to reduce oxygen inclusion as it stuffs the bag. Pressure can be varied to accommodate different feed products. Baggers are available for rent or purchase. Bags come in different sizes and lengths.

Clear a flat space of rocks and debris, and make sure the space is large enough for equipment to move around to fill bags. Be sure to inspect bags and plastic bunker covers for tears or holes because oxygen penetration can cause additional spoilage. Monitor bags and bunkers for evidence of wildlife depredation, and take steps to reduce depredation and silage losses.



**Photo 3. Storing corn grain as high-moisture corn in a bunker silo.** (Greg Lardy, NDSU)



**Photo 4. A large-capacity bagger for placing feed ingredients in anaerobic storage.** (Greg Lardy, NDSU)

## Oxygen-limiting Silos

Whole high-moisture grain may be stored in certain types of oxygen-limiting silos. One advantage of these structures is that corn can be stored at lower moisture levels (22% to 26% moisture) than in bunker or plastic bag storage structures.

Check with the manufacturer to be sure your silo is designed to handle whole high-moisture grain prior to placing grain in the structure.

## Additives

Certain additives may be beneficial in reducing fermentation and feedout losses in HMC. Lactic acid bacteria inoculants that have been developed specifically for high-moisture corn should be considered when the corn has been frozen or is harvested below the optimal moisture content. The use of acid preservatives may be warranted if substantial ear mold damage has occurred and mycotoxin production is prevalent in the standing crop. When considering the use of an additive for high-moisture corn, be sure to ask the manufacturer for research data to support efficacy claims.

## Nutrient Content and Feeding Recommendations

The energy content of high-moisture corn is similar to dry-rolled corn (**Table 1**). However, high-moisture corn typically has higher ruminal and total tract digestion than dry-rolled corn. The higher ruminal digestion rate means producers need to pay particular attention to bunk management when feeding high-moisture corn.

Proper bunk management includes taking care to thoroughly mix all ingredients in the ration, including ionophores or other supplements, feeding at the same time each day, adjusting feed delivered to intake of the animals and observing cattle for signs of acidosis or feed refusal.

In addition, because of its moisture content, high-moisture corn is prone to more rapid deterioration in the bunk than dry-rolled grain. Increased spoilage losses can occur during storage, especially during the

summer, if the high-moisture grain is not properly packed, covered and managed.

Research in Nebraska has documented advantages of feeding mixtures of high-moisture corn and dry-rolled corn in the finishing period. Other research has shown advantages of feeding mixtures of high-moisture corn and wet distillers grain during the finishing period as well.

### Immature Corn

Immature corn may be harvested effectively as high-moisture corn for feed. Research at NDSU indicates light-test-weight, immature corn harvested and fed as high-moisture grain produced equal performance as mature corn grain in feedlot steers.

Analyze grain for nutrient content to more accurately formulate rations because light-test-weight corn often contains more protein than heavy corn.

## Pricing High-moisture Corn

When pricing corn, take into account differences in moisture level, particularly with high-moisture corn. Dry corn typically is traded at 15.5% moisture. High-moisture corn can have moisture levels from 24% to 33%, which necessitates adjustment to a moisture level. Also consider drying costs in pricing wet corn.

### Summary

For beef cattle and dairy producers, harvesting corn as high-moisture grain offers many advantages. Nutrient content and moisture of high-moisture corn may vary depending on harvest timing, yield, variety and other factors. To reduce storage and spoilage losses, follow proper processing and storage recommendations with particular attention to kernel processing, oxygen exclusion/proper packing and covering bunker silos to reduce spoilage.

Take care when feeding high-moisture corn because it does have a faster ruminal digestion rate than dry corn. Consequently, better bunk management typically is required for successful feeding.

**Table 1. Nutrient content of different corn harvest, storage or processing methods.**

Corn Type	Dry Matter	TDN	NE <sub>m</sub>	NE <sub>g</sub>	CP
		%	Mcal/lb	Mcal/lb	%
<b>Grain</b>					
Dry-rolled Corn	86	90	1.02	0.70	9.8
Ear Corn	87	83	0.92	0.62	9.0
High-moisture Corn	75	90	1.02	0.70	10.0

*"Alternative Feeds for Ruminants" (NDSU AS1182)*

<https://www.ndsu.edu/agriculture/extension/publications/alternative-feeds-ruminants>

This publication was originally authored by Greg Lardy, NDSU Animal Sciences Department Head, and Vern Anderson, former NDSU animal scientist.

NDSU Extension does not endorse commercial products or companies even though reference may be made to tradenames, trademarks or service names.

**For more information on this and other topics, see [www.ndsu.edu/extension](http://www.ndsu.edu/extension)**

County commissions, North Dakota State University and U.S. Department of Agriculture cooperating. NDSU does not discriminate in its programs and activities on the basis of age, color, gender expression/identity, genetic information, marital status, national origin, participation in lawful off-campus activity, physical or mental disability, pregnancy, public assistance status, race, religion, sex, sexual orientation, spousal relationship to current employee, or veteran status, as applicable. Direct inquiries to Vice Provost, Title IX/ADA Coordinator, Old Main 100, 701-231-7708, [ndsu.eoaa@ndsu.edu](mailto:ndsu.eoaa@ndsu.edu). This publication will be made available in alternative formats for people with disabilities upon request, 701-231-7881.

2M-7-10; web-1-16; web-10-24