

Name: \_\_\_\_\_

**Quiz 4 Solutions**

Please write your solutions to the following exercises in the space provided. You should write legibly and fully explain your work.

*Good Luck!*

(1) Let  $A_1 = \begin{bmatrix} 1 & 0 \\ 0 & 2 \end{bmatrix}$ ,  $A_2 = \begin{bmatrix} 0 & 1 \\ 2 & 4 \end{bmatrix}$ ,  $A_3 = \begin{bmatrix} 2 & 5 \\ 10 & 24 \end{bmatrix}$  and  $B = \begin{bmatrix} -1 & 3 \\ 6 & 10 \end{bmatrix}$ .

Write  $B$  as a linear combination of  $A_1$ ,  $A_2$  and  $A_3$  by solving a system of linear equations. [10 pts]

*Solution:* We need to find scalars  $c_1, c_2, c_3 \in \mathbb{R}$  such that

$$B = \begin{bmatrix} -1 & 3 \\ 6 & 10 \end{bmatrix} = c_1 A_1 + c_2 A_2 + c_3 A_3 = \begin{bmatrix} c_1 + 2c_3 & c_2 + 5c_3 \\ 2c_2 + 10c_3 & 2c_1 + 4c_2 + 24c_3 \end{bmatrix}.$$

Thus, we need to solve the system

$$\begin{aligned} c_1 + 2c_3 &= -1 \\ c_2 + 5c_3 &= 3 \\ 2c_2 + 10c_3 &= 6 \\ 2c_1 + 4c_2 + 24c_3 &= 10 \end{aligned}$$

We row reduce the augmented matrix of this system

$$\left[ \begin{array}{ccc|c} 1 & 0 & 2 & -1 \\ 0 & 1 & 5 & 3 \\ 0 & 2 & 10 & 6 \\ 2 & 4 & 24 & 10 \end{array} \right] \longrightarrow \left[ \begin{array}{ccc|c} 1 & 0 & 2 & -1 \\ 0 & 1 & 5 & 3 \\ 0 & 2 & 10 & 6 \\ 0 & 4 & 20 & 12 \end{array} \right] \longrightarrow \left[ \begin{array}{ccc|c} 1 & 0 & 2 & -1 \\ 0 & 1 & 5 & 3 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right].$$

We let  $c_3 = t \in \mathbb{R}$ . Then

$$\begin{aligned} c_2 &= -5c_3 + 3 = -5t + 3 \\ c_1 &= -2c_3 - 1 = -2t - 1 \end{aligned}$$

Letting  $t = 1$ , we have

$$B = -3A_1 - 2A_2 + A_3.$$

- (2) Let  $A = \begin{bmatrix} 3 & 0 \\ -1 & 5 \end{bmatrix}$  and  $B = \begin{bmatrix} 4 & -2 & 1 \\ 0 & 2 & 3 \end{bmatrix}$ . Compute  $AB$ . [5 pts]

*Solution:*

$$\begin{bmatrix} 3 & 0 \\ -1 & 5 \end{bmatrix} \begin{bmatrix} 4 & -2 & 1 \\ 0 & 2 & 3 \end{bmatrix} = \begin{bmatrix} 12 & -6 & 3 \\ -4 & 12 & 14 \end{bmatrix}.$$

- (3) Suppose that  $\mathbf{u}, \mathbf{v}$  are linearly independent vectors. Show that the vectors  $\mathbf{u} + \mathbf{v}, \mathbf{u} - \mathbf{v}$  are also linearly independent. [5 pts]

*Solution:* Suppose that for some scalars  $c_1, c_2 \in \mathbb{R}$  we have the equation

$$c_1(\mathbf{u} + \mathbf{v}) + c_2(\mathbf{u} - \mathbf{v}) = \mathbf{0}.$$

Then, rearranging this equation, we see that

$$(c_1 + c_2)\mathbf{u} + (c_1 - c_2)\mathbf{v} = \mathbf{0}.$$

Since  $\mathbf{u}$  and  $\mathbf{v}$  are linearly independent, it must be the case that

$$c_1 + c_2 = 0$$

$$c_1 - c_2 = 0$$

This system implies that  $c_1 = -c_2$  and  $c_1 = c_2$ . However, this can only happen if  $c_1 = c_2 = 0$ . So, by definition,  $\mathbf{u} + \mathbf{v}$  and  $\mathbf{u} - \mathbf{v}$  must also be linearly independent.